

MEDICATIONS which INTERFERE with ALLERGY SKIN TESTS

Certain over-the counter and prescription medications contain ingredients which affect allergy skin tests. Check labels of all medications you are using (including eye drops and nasal sprays) to determine whether your medication contains any ingredient listed below. For questions regarding ingredients contact your pharmacist. The medications listed must be held for at least the amount of time indicated prior to allergy skin tests.

The following **EYEDROPS** must be held at least **7 days prior to tests:**

Pataday, Patanol, Optivar, Zaditor, Alaway, Elestat, olopatadine, azelastine, ketotifen, epinastine Pheniramine (e.g. Visine Allergy Eye drops) must be held for > 48 hours prior to tests

The following **NASAL SPRAYS** must be held for > 7 days prior to tests: **Astelin, Astepro, azelastine, Patanase, olopatidine**

ORAL MEDICATIONS

Minimum time to be held prior to allergy skin tests

Benadryl Diphenhydramine (allergy medications and nonprescription sleep aids)	48 Hours
Doxylamine Pyrilamine Pheniramine (in allergy, cold and sinus preparations)	48 Hours
Phenergan Promethazine (in prescription cough syrups and anti-nausea)	48 Hours
Periactin Cyproheptadine (appetite stimulant and other uses)	48 Hours
Meclizine Dimenhydrinate Antivert Bonine Dramamine (motion sickness)	48 Hours
Ranitidine Famotidine Nizatidine (indigestion, heartburn medications)	48 Hours
Chlor-trimeton Chlorpheniramine (in allergy, cold, and sinus preparations)	5 Days
Tussionex Cough Syrup (contains chlorpheniramine)	5 Days
Zyrtec Xyzal Atarax Vistaril Cetirizine Levocetirizine Hydroxyzine (allergy & itch)	5 Days
Loratadine (in nonprescription allergy medications including Claritin & Alavert)	7 Days
Fexofenadine Allegra Clarinex Desloratadine Brompheniramine Brovex, Loradane	7 Days
Remeron Mirtazapine (treatment of pain, depression, appetite stimulant)	7 Days
Tricyclics (headache, neuralgia, other chronically painful conditions; doxepin also used for itch)	7 Days
Amitriptyline Elavil Imipramine Tofranil	7 Days
Doxepin, Sinequan, Pamelor, Nortriptyline	10 Days

Psychotropic -- **Zyprexa, chlorpromazine, Thorazine, perphenazine, Clozaril clozapine** may affect allergy tests . These medications should not be modified except by instruction of the prescribing physician. Please notify the clinic in advance of your appointment if you are taking these medications.

If you are taking medication with antihistamine effects which cannot be stopped because of the severity of your condition or any other reason, continue taking the medication and let the clinic know in advance of your initial visit. In most cases, alternative medications which will not affect skin tests can be provided at the time of your initial clinic visit with skin testing postponed for a subsequent appointment.

Medications indicated below **DO NOT** affect allergy skin tests and **MAY BE CONTINUED**

Allergy medications

Pseudoephedrine, Phenylephrine Decongestants
 Singulair
 Prednisone, Prednisolone, Methylprednisolone, Medrol
 Lotemax Eyedrops
 Fluticasone, Flonase, Veramyst, Nasonex, Nasocort Aq., Rhinocort, Omnaris Nasal Sprays Mucinex,
 Guiafenesin

All asthma inhalers

Dextromethorphan Cough Medications (E.G. Delsym)
 Prescription Stomach Acid Reducers (Protonix, Prevacid, Nexium, Aciphex) Omeprazole, Prilosec
 Other medications without effect on allergy skin testing include antibiotics, high blood pressure, arthritis, cholesterol, diabetes, cardiac, anticonvulsant, and thyroid medications, most insomnia, anxiety, depression medications including Ambien, Lunesta, Xanax, Ativan, Valium, Prozac, Effexor, Zoloft, bupropion, Wellbutrin, fluoxetine, sertraline, Celexa, Lexapro, and eye drops used for glaucoma.

MEDICAMENTOS que INTERFEREN con las PRUEBAS DE ALERGIAS DE LA PIEL

Ciertos medicamentos de venta libre y recetados contienen ingredientes que afectan las pruebas de alergias de la piel.

Revise las etiquetas de todos los medicamentos que está usando (incluyendo las gotas para los ojos y los aerosoles nasales) para determinar si su medicación contiene algún ingrediente a continuación. Si tiene preguntas sobre los ingredientes, póngase en contacto con su farmacéutico. Los medicamentos listados deben suspenderse durante al menos la cantidad de tiempo indicada antes de las pruebas de alergias de la piel.

Las siguientes **GOTAS DE OJOS** deben suspenderse al menos **7 días antes de las pruebas:**

Pataday, Patanol, Optivar, Zaditor, Alaway, Elestat, olopatadina, azelastina, ketotifeno, epinastina Feniramina (por ejemplo, gotas Visine Allergy Eye) debe suspenderse durante más de **48 horas antes de las pruebas**

Los siguientes **AEROSOLES NASALES** deben suspenderse por más de **7 días antes de las pruebas:**

Astelin, Astepro, azelastine, Patanase, olopatidine

MEDICAMENTOS ORALES

Tiempo mínimo de suspensión antes de las pruebas de alergia de la piel

Benadryl Difenhidramina (medicamentos para la alergia y ayudas para dormir de venta libre)	48 horas
Doxilamina Pirilamina Feniramina (en preparaciones de alergia, resfriado y sinusal)	48 horas
Fenergan Prometazina (en jarabes para la tos de prescripción y anti-náuseas)	48 horas
Periactin Ciproheptadina (estimulante del apetito y otros usos)	48 horas
Meclizina Dimenhydrinato Antivert Bonine Dramamine (mareo por movimiento)	48 horas
Ranitidina Famotidina Nizatidina (indigestión, medicamentos para la acidez estomacal)	48 horas
Clor-trimeton Clorfenamina (en preparaciones para alergias, resfriados y sinusales)	5 días
Jarabe para la tos Tussionex (contiene clorfenamina)	5 días
Zyrtec Xyzal Atarax Vistaril Cetirizina Levocetirizina Hidroxicina (alergia y picazón)	5 días
Loratadina (en medicamentos para la alergia sin receta incluyendo Claritin & Alavert)	7 días
Fexofenadina Allegra Clarinex Desloratadina Bromfeniramina Brovex, Lorlane	7 días
Remeron Mirtazapina (tratamiento del dolor, depresión, estimulante del apetito)	7 días
Tricíclicos (dolor de cabeza, neuralgia, otras condiciones crónicamente dolorosas; doxepina también se usa para la picazón)	
Amitriptilina Elavil Imipramina Tofranil	7 días
Doxepina, Sinequan, Pamelor, Nortriptilina	10 días

Psicotrópico: **Zyprexa, clorpromazina, Toracina, Perfenazina, Clozaril clozapina** puede afectar las pruebas de la alergia. Estos medicamentos no deben modificarse excepto por instrucción del médico que prescribe. Por favor, notifique a la clínica con antelación de su cita si está tomando estos medicamentos.

Si está tomando medicamentos con efectos antihistamínicos que no se pueden detener debido a la gravedad de su afección o cualquier otra razón, continúe tomando el medicamento e informe a la clínica con anticipación a su visita inicial. En la mayoría de los casos, se pueden proporcionar medicamentos alternativos que no afecten las pruebas en el momento de su visita inicial a la clínica con las pruebas dermatológicas pospuestas para una cita posterior.

Los medicamentos indicados a continuación **no afectan las pruebas de alergias de la piel y PUEDEN SER CONTINUADAS**

Medicamentos para la alergia:

Pseudoefedrina, descongestionantes de fenilefrina

Singulair

Prednisona, Prednisolona, Metilprednisolona, Medrol

Gotas de ojos Lotemax

Aerosoles nasales Fluticasona, Flonase, Veramyst, Nasonex, Nasocort Aq., Rhinocort, Omnis

Mucinex, Guiafenesin

Todos los inhaladores para el asma

Medicamentos para la tos de dextrometorfano (p. ej., Delsym)

Reductores de ácido estomacal recetados (Protonix, Prevacid, Nexium, Aciphex)

Omeprazol, Prilosec

Otros medicamentos sin efecto en las pruebas de alergias de la piel incluyen antibióticos, medicamentos para la presión arterial alta, artritis, diabetes con colesterol, medicamentos cardíacos, anticonvulsivos y para la tiroides, la mayoría de los medicamentos para el insomnio, la ansiedad y la depresión, incluidos Ambien, Lunesta, Xanax, Ativan, Valium, Prozac, Effexor, Zoloft, bupropión, Wellbutrin, fluoxetina, sertralina, Celexa, Lexapro y gotas para los ojos para el glaucoma

Allergy Testing

Patient Information

Preparation for Allergy Test

DON'TS

Do not take Claritin, Clarinex, Zyrtec, Allergra or any other allergy medications for 5-7 days prior to the allergy test.

Do not take over the counter antihistamines (cold & sinus medications, sleep aids like Tylenol. PM, Benadryl, Dimetapp, Alevert, Travist, Chlor-Trimeton) 3-4 days before the test.

Do not take medications such as Tagament, Pepcid, or. Zantac 1 day prior to testing, as these contain antihistamine.

Do not wear cologne or scented body lotion. Other allergy patients could be sensitive. Deodorant is fine.

Do not use topical corticosteroids 21 days prior to skin testing.

Do not take a flu shot or any other immunizations 3 days before allergy testing.

TAKING ANY OF THE ABOVE MEDICATIONS CAN ALTER YOUR ALLERGY TEST! IF YOU ARE UNSURE ABOUT A MEDICATION, PLEASE ASK THE NURSE PRIOR TO YOUR APPOINTMENT.

DO'S

Do dress accordingly. If the test is being performed on the skin of your back, you must be prepared to lay still, face down, with your back exposed for 15 minutes. If the test is being performed on the arms, you will need to have your forearms and biceps exposed (both arms) for 15 minutes.

Your practitioner will review your results and discuss with you the appropriate treatments options. Please resume all necessary medications following the test.



Medications to Avoid Before Allergy Skin Testing

Suppressant Effects of Drugs on Immediate Skin Tests

Medications	Mean Days Suppressed	Max Days Suppressed
First Generation Antihistamines	2	5
Second Generation Antihistamines	2	7
Antihistamine Nasal Sprays	0	1
Antihistamine Eye Drops	0	1
Tricyclic Antidepressants and Tranquilizers		14
Histamine2 Antihistamines (H2 Blocker)	0	2
Topical Corticosteroids		Up to 21

Medications that DO NOT Need to be Stopped Prior to Allergy Skin Prick Testing

Angiotensin-Converting Enzyme (ACE) Inhibitors	Nasal Steroid Sprays
<ul style="list-style-type: none"> • Benazepril • Captopril • Enalapril • Lisinopril • Perindopril • Quinapril • Ramipril 	<ul style="list-style-type: none"> • Beclomethasone Dipropionate Nasal • Budesonide Nasal • Ciclesonide Nasal • Fluticasone Propionate • Fluticasone Furoate Nasal • Fluticasone Furoate Nasal • Oymetazoline • Triamcinolone Acetonide
Immunosuppressant	Norepinephrine Reuptake Inhibitors (SNRIs)
<ul style="list-style-type: none"> • Cyclosporin 	<ul style="list-style-type: none"> • Duloxetine • Venlafaxine
Protein Pump Inhibitors (PPIs)	Serotonin Reuptake Inhibitors (SNRIs)
<ul style="list-style-type: none"> • Esomeprazole • Lansoprazole • Omeprazole • Pantoprazole • Rabeprazole 	<ul style="list-style-type: none"> • Citalopram • Escitalopram • Fluoxetine • Paroxetine • Sertraline